

AUTHOR INDEX

Bold type indicates authors of commentaries, editorials, and Resource Corners; (No.) indicates issue number.

- Aaron SD, 12
Aguilar C, 47
Allen C, 91
Allen IE, 69
Alpers JH, 51
Amiel SA, 90
Amin S, 57
Anderson I, 55
Ankum WM, 117
Arroyo V, 20
ASA and Carotid Endarterectomy (ACE) Trial
Collaborators, 10
Ascherio A, 115
Aschwanden M, 108
Assendelft WJ, 62
ATLANTIC Investigators, 85
Atmar RL, 92
Atorvastatin versus Revascularization
Treatment Investigators, 7
Attia J, 32, 33
Ausejo M, 50
- Bakx JC, 37
Baldwin D, 103
Barnett HJ, 10
Barron HV, 36
Bates ER, 36, 81
Bath PMW, 11
Bauer DC, 33
Bauer M, 105
Bauer TT, 89
Bell-Syer S, 99
Benavente O, 86
Bender BS, 49
Bent S, 48
Berger PB, 71
Berman B, 23
Besag FM, 22
Bijlsma JW, 62
Birkmeyer JD, 72
Black DM, 58
Bobes J, 103
Bollini P, 25
Booth A, A-16 (No. 3)
Borzak S, 42
Bota GW, 13
Bouma M, 113
Bradley CP, 64
Bradley KA, 67
Bronson DL, 95
Brott BC, 84-85
Brown G, 102
Brown WV, 7
Browner WS, A-18 (No. 3)
Buckley RG, 117
Burkhoff D, 85
Burroughs AK, 68
Buttery J, 27
Byles JE, 74
- Califf RM, 5
Canadian Task Force on Preventive Health
Care, 107, 112
Carey TS, 26
Carolei A, 114
Carpenter S, 35
Chambers GK, 1
Cherkin DC, 23
Chey WD, 34, 35
Christensen K, 29
Cleland J, 4
Collins D, 44
Collins JS, 46
Cometta A, 53
Cook DJ, 32, 89, 111
Cooper BA, 73
Cooper KG, 28
Coughlan JL, 15
Cowan J, A-19 (No. 3)
Crews FT, 26
Cullum N, 59
- Dales RE, 12
Daley J, 72
Dammers JW, 56
Datta M, 45
Davey Smith G, 43
Davidson RA, 37
Davies MJ, 17
Davies RJ, 16
Davis D, 75
de Bock R, 53
de Boer MJ, 81
de Sonnaville JJ, 113
Deasy JM, 60
DECODE Study Group, 70
Deeks J, 59
Dekker J, 62
Dekker JH, 113
Del Mar C, 91, 101
Delaney B, 34-35
Delis KT, 109
Department of Veterans Affairs Cooperative
Study Group, 14
Detsky AS, 76
Deupree RH, 14
Dignam J, 54
Disney JD, 117
Donnelly S, 98
Donson H, 102
Döpfner S, 105
Douglas N, 69
Douketis J, 108-109
Drakulovic MB, 89
Durham SR, 65
Dusseldorp E, 87
- Ebrahim S, 43
Eikelboom JW, 73
- Eisenberg PR, 71**
Ekblom T, 95
Ellis SG, 71
Erblund ML, 14
Eslick GD, 74
Etchells E, A-15 (No. 3)
Ettinger B, 58
European Diabetes Epidemiology Group, 70
Evaluation of Platelet IIb/IIIa Inhibition in
Stenting Investigators, 5
Evans AT, 113
Every NR, 36
- Fabris L, 13
Fahey T, 48
Faure M, 103
Fekete T, A-18 (No. 3)
Felson DT, 57
Fenlon HM, 110
Fiset V, 66
Fisher B, 54
Fitzpatrick R, 106
Fleming DM, 93
Fletcher AW, 59
Fletcher RH, 110
Forster J, 50
FRagmin and Fast Revascularisation during
InStability in Coronary artery disease
(FRISC II) Investigators, 82, 83
Frankel M, 11
Frazier OH, 84
Freemantle N, 4, 75
Freifeld A, 52
Fuller P, 17
- Ganda OP, 70**
Garbutt JC, 26
Garratt AM, 28
Gehrig A, 108
Genest J Jr, 73
Gherardi G, 18
Gibson JN, 100
Gibson PG, 14, 15
Gillespie LD, 63
Gillespie WJ, 63, 100
GISSI-Prevenzione Investigators, 6
Glanville J, A-11-13 (No. 3)
Glasziou P, A-11-14 (No. 1), 91
Gluckman R, 43, 87
Goldberg AS, 30
Golish JA, 16
Goodwin GM, 25
Goodwin PJ, 106
Gorman JD, 117
Goulis J, 68
Grady D, 48
Graham DY, 97
Grant AD, 45

- Grant AM, 28
Grant IC, 100
Green A, 101
Gross M, 29
Grunfeld E, 106
GUSTO-II Investigators, 71
Guyatt G, 33
Guyatt GH, 111
Hankey G, 115
Hansson L, 95
Harper S, 96
Harrison J, 4
Harrison KJ, 69
Hart R, 9, 86
Hatala R, 32
Hatlebakk JG, 64
Havranek EP, 44
Hayden FG, 92
**Haynes B, A-11 (No. 1), A-11-14 (No. 1)
10, A-13 (No. 2)**
Healy D, 28
Heaney A, 46
Heard AR, 51
Heart Outcomes Prevention Evaluation Study
Investigators, 41
Heiligenstein J, 55
Heller SR, 90
Henderson M, 2
Henderson W, 72
Henry D, 93, 96-97
Henry RL, 15
Hickam DH, 8
Hillegass WB, 5
Hochman JS, 42
Holm HA, 75
Holmes DR Jr, 71
Hoorntje JC, 81
Horowitz JD, 88
Horvath KA, 84
Hulm S, 30
Huskinson EC, 98
Hyggen A, 64
- International Antimicrobial Therapy
Cooperative Group of the European
Organization for Research and
Treatment of Cancer, 53
Ishani A, 94
- Jaeger KA, 108
Jeschke R, A-16 (No. 2)
Jamal SA, 57
Jeanneret C, 108
Jenkinson C, 16
Johnson ES, 9
Joshiyura KJ, 115
Juul P, 29
- Kamath S, 82-83**
Kapral MK, 112
Kern WV, 53
Khuri SF, 72
- Kiel DP, 63
Kimmey MB, 19
King KJ, 117
Klausen JH, 117
Koes BW, 23
Korula J, 20
Kraaij V, 87
Kreder HJ, 98
Kronzon I, 112
Krumholz HM, 36
Kumar M, 21
Kumwenda NI, 116
Kwiatkowski TG, 11
- Labs KH, 108
Lacasse Y, 111
Laine L, 35, 96
Lam RW, 105
Lamba GS, 21
Lanes SF, 9
Lang DM, 15
Langhorne P, 114
Langman MJ, 74
Lao L, 23
LaValley MP, 57
Laville M, 18
Leandro G, 68
Lee CH, 1
Lee TH, 76
Lefebvre C, A-11-13 (No. 3)
Lennox AF, 109
Li JT, 65
Libman RB, 11
Lim LL, 74
Lincoff AM, 5
Lindholm LH, 95
Lip GYH, 82-83
Lipman T, A-16 (No. 1)
Live Attenuated Influenza Virus Vaccine in
Healthy Adults Trial Group, 49
Locke KA, 76
Lohr KN, 26
Lonn E, 73
- Mac Donald R, 94
Madsen PH, 64
Maes S, 87
Magee LA, A-19 (No. 2)
Makela M, 47
Mallon KP, 49
Malmberg K, 41
Mangione CM, 76
Mansell P, 90
Manson JE, 115
Mant D, 106
Marcantonio ER, 76
March RJ, 84
Marchigiani D, 52
Margetts P, 33
Margo K, 99
Marini C, 114
Marley JE, 88
- Maron DJ, 7
Marshall JC, 60
Mårtensson L, 61
Mason J, 4
Massel D, 6
Mayo-Smith M, 26
Mazur LJ, 102
McBride R, 86
McConnachie A, 31
McGettigan P, 96-97
McGinn LK, 104
McKelvie RS, 3, 88
McKenzie SA, 17
McLeod RS, 107
McNutt R, 66
Mendelman PM, 49
MERIT-HF Study Group, 3
Meulman J, 87
Millard PS, 94
Miotto PG, 116
Misra A, 21
Mitlak BH, 58
Moffett JK, 99
Moliterno DJ, 5
Monto AS, 93
Moody KA, 116
Moore AP, 24
Morley C, 31
Mullins R, 16
Mulrow CD, 2, A-19 (No. 2)
Multiple Outcomes of Raloxifene Evaluation
(MORE) Investigators, 58
Murthy NS, 21
Murthy U, 34, 35
Mylotte JM, 77
- National Institute of Neurological Disorders
and Stroke Recombinant Tissue
Plasminogen Activator Stroke
Study Group, 11
National Registry of Myocardial
Infarction 2 Participants, 36
National Research Council Study Group on
Stroke in the Young, 114
Naughton BJ, 77
Navasa M, 20
Neale R, 101
Neville BG, 22
Nichol KL, 49, 92-93
Niewoehner DE, 14, 111
Nikodem C, 61
Norwegian Heartburn Study Group, 64
Nunes DP, 110
- O'Connor AM, 66
Olson DH, 67
Oostendorp RA, 62
O'Rourke A, A-16 (No. 3)
Osetamivir Study Group, 92
- Pampallona S, 25
Parker MJ, 63

- Parkin DE, 28
 Paroxetine Study Group, 103
 Parrino TA, 32
 Parsons L, 36
 Paton JY, 17
 Paykel ES, 104
 Pearce LA, 86
 Perez E, 54, 107
 Perna A, 18
 Pham B, 50
 Phelps KR, 58
 Phillips E, 52-53
 Picciotto A, 17
 Pilotto LS, 13
 Pitt B, 2, 7
 Polmear A, A-15 (No. 2)
 Ponc R, 19
 Priore R, 77
 Pullicino P, 10, 86

 Randomized Aldactone Evaluation Study
 Investigators, 2
 Rao G, 51
 Redmond HP, 60
 Reedy K, 77
 Reginster JY, 62
 Reichen J, 21, 68
 Reis S, 23, 91
 Remme WJ, 2
 Revicki D, 55
 Richards IJ, 51
 Richardson WS, A-17-18 (No. 3)
 Robins SJ, 44
 Rofecoxib Osteoarthritis Endoscopy Study
 Group, 96
 Ross S, 31
 Ross SD, 69
 Rostom A, 66
 Rounding C, 30
 Rowe BH, 12, 13
 Rubins HB, 44
 Ruggenenti P, 18
 Rydén L, 41

 Saenz A, 50
 Saint S, 48
 Saklayen MG, 1
 Saltzman S, 77
 Sarin SK, 21
 Sassan-Morokro M, 45
 Saunders MD, 19
 Scharwächter I, 103
 Schilling M, 92
 Schmidt S, 85
 Schneiderman H, A-18-19 (No. 3)
 Schroy PC 3d, 110

 Schulman SP, 85
 Schulz KF, A-11-12 (No. 2)
 Scott J, 104
 Scott RC, 22
 Serunkuma S, 109
 Shaw S, 34
 Sheldon TA, 59
 Shivkumar K, 4
 Shock Investigators, 42
 Sibai B, A-19 (No. 2)
 Sigal RJ, 90
 Silver FL, 112
 Simms RW, 57
 Simon GE, 55
 Simon LS, 97
 Simon T, 96
 Sleeper LA, 42
 Smith BJ, 13
 Smith DM, 77
 Song F, 59
 Sookhai S, 60
 Sort P, 20
 Sparandero F, 47
 Spinhoven P, 8
 Stein DJ, 103
 Stewart S, 88
 Stiasny K, 24
 Stoodley RG, 12
 STOP-Hypertension-2 study group, 95
 Stradling JR, 16
 Straus S, A-11-14 (No. 1), A-15 (No. 1)

 Taha TE, 116
 Talley NJ, 46, 74
 Tatineni S, 5
 Taylor DW, 10
 Teasdale JD, 104
 Temple LK, 107
 Thomson H, 31
 Thomson O'Brien MA, 75
 Tibaldi G, 25
 Tinnion ON, 27
 Torgerson D, 99
 Torres A, 89
 Toskes P, 35
 Totaro R, 114
 Transmyocardial Carbon Dioxide Laser
 Revascularization Study Group, 84
 Triplehorn C, 94, 116
 Tunick PA, 112

 U.K. Lispro Study Group, 90

 VA National Surgical Quality Improvement
 Program, 72
 Vaccarino V, 36

 van Baar ME, 62
 van den Bosch WJ, 37
 van den Hoogen HJ, 37
 van Elderen T, 87
 van Peski-Oosterbaan AS, 8
 van Rood Y, 8
 van Tulder MW, 23
 van Zanten SV, 19
 Varga E-M, 65
 Veering MM, 56
 Vermeulen M, 56
 Veterans Affairs High-Density Lipoprotein
 Cholesterol Intervention Trial
 Study Group, 44
 Vittinghoff E, 48
 von Dadelnszen P, A-19 (No. 2)

 Waddell G, 100
 Walker SM, 65
 Wallin G, 61
 Walsh T, 52
 Wang EE, 107
 Waters D, 7
 Watson R, 31
 Watson RG, 46
 Weaver AL, 97
 Webb JG, 42
 Weinreb H, 56
 Wentworth CE 3d, 9
 West SL, 26
 Wetter TC, 24
 Wiebe S, 22
 Wiktor SZ, 45
 Willenbring ML, 67
 Williams G, 101
 Williams JW Jr, 47
 Wilson MC, A-17-18 (No. 3)
 Wilson P, 31
 Wilson SA, 103
 Wilt T, 94
 Winkelmann J, 24
 Wolmark N, 54
 Wong E, 111
 Wong JG, 32
 Wright JM, 1

 Xia HH, 74

 Young P, 4
 Yusuf S, 73

 Zannad F, 2
 Zar HJ, 102
 Zeleznik J, 59
 Zijlstra F, 81

A

- Acupuncture
 - Review: The effectiveness of acupuncture for nonspecific low-back pain is unclear, 23
- Adaptation, psychological
 - Review: Psychoeducational programs reduce MI recurrence and improve some physical health outcomes, 87
- Administration, buccal
 - Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy, 22
- Administration, rectal
 - Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy, 22
- Adrenal cortex hormones
 - Review: Vitamin D plus calcium is more effective than no therapy or calcium alone in corticosteroid-induced osteoporosis, 57
- Adrenergic-antagonists
 - Endoscopic ligation was better than propranolol for prevention of bleeding from esophageal varices, 21
- Adrenergic beta-agonists
 - Controlled- and extended-release metoprolol reduced mortality in congestive heart failure, 3
 - Conventional and newer hypertensive drugs had similar efficacy in elderly persons with hypertension, 95
 - Review: β -agonists plus ipratropium bromide improve airflow obstruction in adults with acute asthma, 12
 - Review: Long-term β -blocker use reduces mortality after myocardial infarction, 4
- Age factors
 - Younger women with acute MI had more in-hospital deaths than men of the same age, 36
- Albumins
 - Plasma volume expansion with albumin reduced renal impairment and death in cirrhosis and spontaneous bacterial peritonitis, 20
- Alcoholism
 - Integrated outpatient treatment increased abstinence in men with alcohol-related illness who were ongoing drinkers, 67
 - Review: Studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence, 26
- Aldosterone antagonists
 - Spironolactone reduced mortality in severe congestive heart failure, 2
- Allergens
 - Grass-pollen immunotherapy for 3 to 4 years was still effective 3 years after being discontinued, 65
- Ambulatory care
 - Integrated outpatient treatment increased abstinence in men with alcohol-related illness who were ongoing drinkers, 67
- Amikacin
 - Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 53
- Amoxicillin
 - Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 52,53
 - Review: Penicillin V or amoxicillin is better than placebo and equal to nonpenicillins for acute maxillary sinusitis, 47
- Analgesia, obstetric
 - Cutaneous lumbosacral injections of sterile water were more effective than "placebo" injections for relieving first-stage labor pain, 61
- Angina pectoris
 - Dalteparin reduced death and MI at 1 but not 3 months in unstable coronary artery disease managed noninvasively, 82
 - An invasive strategy reduced the combined rate of death and myocardial infarction in unstable coronary artery disease, 83
 - Transmyocardial revascularization increased exercise duration and reduced angina symptoms, 85
 - Transmyocardial revascularization was effective in refractory angina and left ventricular free-wall ischemia, 84
- Angioplasty, transluminal percutaneous coronary
 - Abciximab and stenting reduced death, myocardial infarction, and repeated revascularization in coronary revascularization, 5
 - Atorvastatin was at least as effective as PTCA for reducing ischemic events in patients with stable coronary artery disease, 7
 - Immediate revascularization for cardiogenic shock after MI reduced 6-month mortality, 42
 - Increased time from assessment to angioplasty was associated with an increased 30-day mortality rate, 71
 - Primary angioplasty had lower long-term mortality than did streptokinase in acute myocardial infarction, 81
- Angiotensin-converting enzyme inhibitors
 - Conventional and newer hypertensive drugs had similar efficacy in elderly persons with hypertension, 95
 - Ramipril reduced mortality and cardiovascular morbidity in high-risk adults, 41
 - Ramipril reduced the rate of progression to end-stage renal disease and overt proteinuria in nondiabetic nephropathy, 18
- Antibiotics
 - Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 52,53
 - Review: Antibiotics have a slight beneficial effect on acute bronchitis, 48
 - Review: Penicillin V or amoxicillin is better than placebo and equal to nonpenicillins for acute maxillary sinusitis, 47
- Antibodies, bacterial
 - Whole-blood antibody tests were not highly sensitive for detecting *Helicobacter pylori* infection, 34
- Anticoagulants
 - Review: Antithrombotic agents prevent stroke in nonvalvular atrial fibrillation, 86
- Antidepressive agents
 - More people with depression continued treatment with fluoxetine than with desipramine or imipramine, 55
 - Review: Lower doses of antidepressant drugs are effective and have fewer adverse effects in depression, 25
- Antihypertensive agents
 - Review: Antihypertensive drugs improve maternal outcomes in mild chronic and pregnancy-induced hypertension (Letter), A-19(No. 2)
 - Review: Low-dose thiazides are the most effective first-line drugs for hypertension, 1
- Anti-inflammatory agents, steroidal
 - Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma, 13

- Methylprednisolone injection improved symptoms for 1 year in patients with the carpal tunnel syndrome, 56
- Antilipemic agents**
- Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high-density lipoprotein cholesterol, 44
- Aspirin**
- Low-dose aspirin reduced combined 3-month risk for stroke, MI, and death after carotid endarterectomy, 10
- Review: Antithrombotic agents prevent stroke in nonvalvular atrial fibrillation, 86
- Review: Aspirin reduces the risk for stroke in patients with previous TIA or stroke but does not have a dose-response effect, 9
- Asthma**
- Attendance at an asthma clinic did not reduce asthma morbidity more than did standard medical treatment, 51
- Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma, 13
- Review: β -agonists plus ipratropium bromide improve airflow obstruction in adults with acute asthma, 12
- Review: Treatment of gastroesophageal reflux does not improve asthma outcomes, 15
- Spacers made from sealed cold-drink bottles were as effective as conventional spacers for children with acute asthma, 102
- Atherosclerosis**
- Review: High levels of homocysteine are associated with an increased risk for cardiovascular disease, 73
- Atrial fibrillation**
- Review: Antithrombotic agents prevent stroke in nonvalvular atrial fibrillation, 86
- B**
- Bacterial infections**
- Plasma volume expansion with albumin reduced renal impairment and death in cirrhosis and spontaneous bacterial peritonitis, 20
- Bed rest**
- Review: Early mobilization may be better than bed rest for medical conditions and after surgery, 91
- Beds**
- Review: Specially designed products to prevent or heal pressure ulcers are more effective than standard mattresses, 59
- Behavior therapy**
- Cognitive behavioral therapy reduced noncardiac chest pain and use of psychological services, 8
- Beta carotene**
- Daily use of β -carotene supplements did not prevent skin cancer, but daily sunscreen use reduced the incidence of squamous-cell carcinomas, 101
- Biopsy, needle**
- Review: Transthoracic needle aspiration biopsy is accurate for diagnosing malignant and specific benign pulmonary lesions, 111
- Blood glucose**
- Insulin lispro reduced nocturnal hypoglycemic episodes in type 1 diabetes mellitus when used to achieve tight glycemic control in an intensified insulin therapy regimen, 90
- A 2-hour oral glucose tolerance test result ≥ 11.1 mmol/L predicted all-cause mortality better than a fasting glucose level ≥ 7.0 mmol/L, 70
- Blood pressure**
- Diastolic blood pressure and weight gain predicted future risk for hypertension, 37
- Brain ischemia**
- Fruit and vegetable intake decreased risk for ischemic stroke, 115
- Long-term survival decreased in young adults after stroke or transient ischemic attack, 114
- Breast feeding**
- HIV transmission risk during breast-feeding was greatest during the early months of life, 116
- Breast neoplasms**
- Review: Routine follow-up tests do not improve survival and quality of life in women after treatment for early-stage breast cancer, 106
- Tamoxifen added to lumpectomy and radiation therapy reduced breast cancer events in ductal carcinoma in situ, 54
- Women with breast cancer were more satisfied with general practitioner care than with outpatient clinic care, 106
- Bronchitis**
- Review: Antibiotics have a slight beneficial effect on acute bronchitis, 48
- Budesonide**
- Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma, 13
- C**
- Calcium channel blockers**
- Conventional and newer hypertensive drugs had similar efficacy in elderly persons with hypertension, 95
- Carbon isotopes**
- The ^{13}C -urea blood test is accurate for detecting *Helicobacter pylori* infection, 35
- Cardiovascular agents**
- Transmyocardial revascularization increased exercise duration and reduced angina symptoms, 85
- Transmyocardial revascularization was effective in refractory angina and left ventricular free-wall ischemia, 84
- Cardiovascular disease**
- Ramipril reduced mortality and cardiovascular morbidity in high-risk adults, 41
- Review: High levels of homocysteine are associated with an increased risk for cardiovascular disease, 73
- Carpal tunnel syndrome**
- Methylprednisolone injection improved symptoms for 1 year in patients with the carpal tunnel syndrome, 56
- Cefotaxime**
- Plasma volume expansion with albumin reduced renal impairment and death in cirrhosis and spontaneous bacterial peritonitis, 20
- Ceftazidime**
- Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 52
- Cerebral ischemia**
- Tissue plasminogen activator improved function at 6 and 12 months after ischemic stroke, 11

Cerebrovascular disorders

- Long-term survival decreased in young adults after stroke or transient ischemic attack, 114
- Low-dose aspirin reduced combined 3-month risk for stroke, MI, and death after carotid endarterectomy, 10
- Review: Aspirin reduces the risk for stroke in patients with previous TIA or stroke but does not have a dose-response effect, 9
- Review: Transthoracic echocardiography and transesophageal echocardiography detect cardiac masses in patients with stroke, 112

Chest pain

- Cognitive behavioral therapy reduced noncardiac chest pain and use of psychological services, 8

Child welfare

- The *Baby Check* booklet did not affect frequency of health service use in infants ≤ 6 months of age, 31

Choice behavior

- Review: Health care decision aids improve knowledge, decrease decisional conflict, and increase active participation, 66

Cisapride

- Omeprazole was better than cisapride or placebo for controlling heartburn at 4 and 8 weeks, 64

Clinical competence

- Review: Interactive, but not didactic, continuing medical education is effective in changing physician performance, 75

Cognitive therapy

- Antidepressants plus cognitive therapy reduced relapse in residual depression, 104
- Cognitive behavioral therapy reduced noncardiac chest pain and use of psychological services, 8

Coin lesion, pulmonary

- Review: Transthoracic needle aspiration biopsy is accurate for diagnosing malignant and specific benign pulmonary lesions, 111

Colonic disorders

- Neostigmine improved acute colonic pseudo-obstruction quickly and effectively, 19
- Virtual colonoscopy was sensitive and specific for detecting colorectal polyps and cancer, 110

Colonoscopy

- Virtual colonoscopy was sensitive and specific for detecting colorectal polyps and cancer, 110

Coronary disease

- Abciximab and stenting reduced death, myocardial infarction, and repeated revascularization in coronary revascularization, 5
- Atorvastatin was at least as effective as PTCA for reducing ischemic events in patients with stable coronary artery disease, 7
- Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high-density lipoprotein cholesterol, 44
- Review: Multiple interventions modestly reduce cardiovascular risk factors in primary prevention, but effects on mortality are uncertain, 43
- Review: Psychoeducational programs reduce MI recurrence and improve some physical health outcomes, 87
- A simple index predicted cardiac complications in elective major noncardiac surgery, 76

Cough

- A 14-day regimen of high-dose inhaled corticosteroids was more effective than placebo for persistent nocturnal cough, 17

Counseling

- Review: Multiple interventions modestly reduce cardiovascular risk factors in primary prevention, but effects on mortality are uncertain, 43

Critical care

- A semirecumbent body position produced a lower rate of nosocomial pneumonia than did a supine position in mechanically ventilated adults, 89

Croup

- Review: Glucocorticoids improve symptoms of croup within 6 hours, 50

Cyclooxygenase inhibitors

- Celecoxib was similar to naproxen for rheumatoid arthritis with fewer endoscopic ulcers, 97
- Rofecoxib caused fewer endoscopic gastroduodenal ulcers than ibuprofen in osteoarthritis, 96

D

Danazol

- Transcervical resection of the endometrium was effective and acceptable at up to 2 years for heavy menstrual bleeding, 28

Databases, bibliographic

- Ovid (Resource Corner), A-15(No. 3)
- PubMed and SUMSearch (Resource Corner), A-16 (No. 3)

Decision support techniques

- Review: Health care decision aids improve knowledge, decrease decisional conflict, and increase active participation, 66

Decubitus ulcer

- Review: Specially designed products to prevent or heal pressure ulcers are more effective than standard mattresses, 59

Depression

- Antidepressants plus cognitive therapy reduced relapse in residual depression, 104
- More people with depression continued treatment with fluoxetine than with desipramine or imipramine, 55
- Review: Lithium augmentation increases treatment response in refractory depression, 105

Diabetes mellitus

- A fasting plasma glucose level ≥ 7.8 mmol/L was sensitive but not specific for identifying poor glycemic control, 113
- Insulin lispro reduced nocturnal hypoglycemic episodes in type 1 diabetes mellitus when used to achieve tight glycemic control in an intensified insulin therapy regimen, 90
- A 2-hour oral glucose tolerance test result ≥ 11.1 mmol/L predicted all-cause mortality better than a fasting glucose level ≥ 7.0 mmol/L, 70

Diagnostic techniques and procedures

- Diagnostic Strategies for Common Medical Problems* (Resource Corner), A-15(No. 1)

Diazepam

- Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy, 22

- Diet
Fruit and vegetable intake decreased risk for ischemic stroke, 115
- Disease transmission, vertical
HIV transmission risk during breast-feeding was greatest during the early months of life, 116
- Disease-free survival
Long-term survival decreased in young adults after stroke or transient ischemic attack, 114
- Diskectomy
Review: Evidence supports surgery for lumbar disc prolapse but is insufficient for degenerative lumbar spondylosis, 100
- Disulfiram
Review: Studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence, 26
- Diuretics
Conventional and newer hypertensive drugs had similar efficacy in elderly persons with hypertension, 95
- Dose-response relationship, drug
Review: Aspirin reduces the risk for stroke in patients with previous TIA or stroke but does not have a dose-response effect, 9
Review: Lower doses of antidepressant drugs are effective and have fewer adverse effects in depression, 25
- Double-blind method
Assessing allocation concealment and blinding in randomized controlled trials: why bother? (Editorial), A-11(No. 2)
- Drug resistance
Review: Lithium augmentation increases treatment response in refractory depression, 105
- Dyspepsia
Empirical eradication therapy was better than endoscopy-based management for *Helicobacter pylori* infection and dyspepsia, 46
- E**
- Echocardiography
Review: Transthoracic echocardiography and transesophageal echocardiography detect cardiac masses in patients with stroke, 112
- Ectopic pregnancy
The high-risk criteria of a clinical prediction model were specific but not sensitive for predicting ectopic pregnancy, 117
- Education, medical
Drug dependence in a journal club (Letter), A-17(No. 3)
Review: Interactive, but not didactic, continuing medical education is effective in changing physician performance, 75
- Emergency treatment
Review: β -agonists plus ipratropium bromide improve airflow obstruction in adults with acute asthma, 12
- Endarterectomy, carotid
Low-dose aspirin reduced combined 3-month risk for stroke, MI, and death after carotid endarterectomy, 10
- Endoscopy
Empirical eradication therapy was better than endoscopy-based management for *Helicobacter pylori* infection and dyspepsia, 46
- Epilepsy
Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy, 22
- Esophageal and gastric varices
Endoscopic ligation was better than propranolol for prevention of bleeding from esophageal varices, 21
- Estrogen antagonists
Transcervical resection of the endometrium was effective and acceptable at up to 2 years for heavy menstrual bleeding, 28
- Estrogens
Raloxifene reduced vertebral fractures in postmenopausal women, 58
- Evidence-based medicine
Advances in evidence-based information resources for clinical practice (Editorial), A-11(No. 1)
Clinical Evidence (Resource Corner), A-15(No. 2)
Diagnostic Strategies for Common Medical Problems (Resource Corner), A-15(No. 1)
Drug dependence in a journal club (Letter), A-17(No. 3)
Incorporating allocation concealment and blinding in randomized controlled trials (Editorial), A-13(No. 2)
Netting the Evidence (Resource Corner), A-16(No. 1)
Ovid (Resource Corner), A-15(No. 3)
PubMed and SUMSearch (Resource Corner), A-16 (No. 3)
UpToDate (Resource Corner), A-16(No. 2)
- Exercise therapy
An 8-session exercise program was effective for subacute or chronic low-back pain, 99
Review: Exercise therapy may reduce pain and disability in osteoarthritis of the hip or knee, 62
Review: Psychoeducational programs reduce MI recurrence and improve some physical health outcomes, 87
- F**
- Family practice
The *Baby Check* booklet did not affect frequency of health service use in infants ≤ 6 months of age, 31
- Fatty acids, omega-3
n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction, 6
- Fenoterol
Spacers made from sealed cold-drink bottles were as effective as conventional spacers for children with acute asthma, 102
- Fever
Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 52,53
- Fibrin fibrinogen degradation products
A negative D-dimer result and low-risk clinical status effectively ruled out DVT in symptomatic patients, 108
A negative D-dimer test result alone or combined with low-risk clinical status effectively ruled out symptomatic DVT, 109
- Fibrinolytic agents
Primary angioplasty had lower long-term mortality than did streptokinase in acute myocardial infarction, 81
Tissue plasminogen activator improved function at 6 and 12 months after ischemic stroke, 11
- Fish oils
n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction, 6
- Fluoxetine
More people with depression continued treatment with fluoxetine than with desipramine or imipramine, 55
- Fluticasone
A 14-day regimen of high-dose inhaled corticosteroids was more effective than placebo for persistent nocturnal cough, 17

- Fruit
Fruit and vegetable intake decreased risk for ischemic stroke, 115

G

- Gastric mucosa
Celecoxib was similar to naproxen for rheumatoid arthritis with fewer endoscopic ulcers, 97
- Gastric mucosa
Rofecoxib caused fewer endoscopic gastroduodenal ulcers than ibuprofen in osteoarthritis, 96
- Gastrointestinal disorders
Endoscopic ligation was better than propranolol for prevention of bleeding from esophageal varices, 21
Review: Treatment of gastroesophageal reflux does not improve asthma outcomes, 15
- Gemfibrozil
Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high-density lipoprotein cholesterol, 44
- Geriatric assessment
Organ system function, general health, social support, and age predicted discharge from a postacute geriatric unit to the community, 77
- Glucocorticoids
Glucocorticoids reduced short-term treatment failure in exacerbations of chronic obstructive pulmonary disease, 14
Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma, 13
Review: Glucocorticoids improve symptoms of croup within 6 hours, 50
- Glucose tolerance test
A fasting plasma glucose level ≥ 7.8 mmol/L was sensitive but not specific for identifying poor glycemic control, 113

H

- Hayfever
Grass-pollen immunotherapy for 3 to 4 years was still effective 3 years after being discontinued, 65
- Health education
Review: Psychoeducational programs reduce MI recurrence and improve some physical health outcomes, 87
- Heart failure, congestive
Controlled- and extended-release metoprolol reduced mortality in congestive heart failure, 3
A multidisciplinary, home-based intervention reduced deaths and readmissions in patients with chronic congestive heart failure, 88
Spironolactone reduced mortality in severe congestive heart failure, 2
- Heartburn
Omeprazole was better than cisapride or placebo for controlling heartburn at 4 and 8 weeks, 64
- Helicobacter* infections
Empirical eradication therapy was better than endoscopy-based management for *Helicobacter pylori* infection and dyspepsia, 46
Review: Pooled heterogeneous studies show an association between *Helicobacter pylori* infection and gastric cancer, 74
The ^{13}C -urea blood test is accurate for detecting *Helicobacter pylori* infection, 35

- Whole-blood antibody tests were not highly sensitive for detecting *Helicobacter pylori* infection, 34
- Hematologic tests
The ^{13}C -urea blood test is accurate for detecting *Helicobacter pylori* infection, 35
- Hemoglobin A, glycosylated
A fasting plasma glucose level ≥ 7.8 mmol/L was sensitive but not specific for identifying poor glycemic control, 113
- Heptanoic acids
Atorvastatin was at least as effective as PTCA for reducing ischemic events in patients with stable coronary artery disease, 7
- Hernia, inguinal
Laparoscopic hernia repair reduced recovery time and days of postoperative analgesia use better than Shouldice repair, 29
- Hip fractures
Review: External hip protectors reduce the risk for hip fractures in elderly persons, 63
- HIV infections
HIV transmission risk during breast-feeding was greatest during the early months of life, 116
Trimethoprim-sulfamethoxazole decreased morbidity and mortality in HIV-1-infected patients with tuberculosis, 45
- Home care services
A multidisciplinary, home-based intervention reduced deaths and readmissions in patients with chronic congestive heart failure, 88
- Homocysteine
Review: High levels of homocysteine are associated with an increased risk for cardiovascular disease, 73
- Hospital mortality
Younger women with acute MI had more in-hospital deaths than men of the same age, 36
Surgical volume was not related to 30-day mortality in 8 common operations, 72
- Hyaluronic acid
Hyaluronic acid improved walking ability and short-term knee function in osteoarthritis of the knee, 98
- Hyperglycemia
A 2-hour oral glucose tolerance test result ≥ 11.1 mmol/L predicted all-cause mortality better than a fasting glucose level ≥ 7.0 mmol/L, 70
- Hyperkalemia
Spironolactone reduced mortality in severe congestive heart failure, 2
- Hypertension
Conventional and newer hypertensive drugs had similar efficacy in elderly persons with hypertension, 95
Diastolic blood pressure and weight gain predicted future risk for hypertension, 37
Review: Antihypertensive drugs improve maternal outcomes in mild chronic and pregnancy-induced hypertension (Letter), A-19(No. 2)
Review: Low-dose thiazides are the most effective first-line drugs for hypertension, 1
- Hypertension, renal
Ramipril reduced the rate of progression to end-stage renal disease and overt proteinuria in nondiabetic nephropathy, 18
- Hypoglycemia
Insulin lispro reduced nocturnal hypoglycemic episodes in type 1 diabetes mellitus when used to achieve tight glycemic control in an intensified insulin therapy regimen, 90

Hypolipoproteinemia

- Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high-density lipoprotein cholesterol, 44

I

Ibuprofen

- Celecoxib was similar to naproxen for rheumatoid arthritis with fewer endoscopic ulcers, 97
- Identifying systematic reviews: key resources (Editorial), A-11(No. 3)
- Rofecoxib caused fewer endoscopic gastroduodenal ulcers than ibuprofen in osteoarthritis, 96

Immunotherapy

- Grass-pollen immunotherapy for 3 to 4 years was still effective 3 years after being discontinued, 65

Infant, newborn

- The *Baby Check* booklet did not affect frequency of health service use in infants ≤ 6 months of age, 31

Influenza

- Intranasal, live, attenuated influenza virus vaccine reduced severe febrile illness in healthy adults, 49
- Oseltamivir once or twice daily safely prevented influenza, 92
- Zanamivir speeds symptom relief in influenza, 93

Injections

- Cutaneous lumbosacral injections of sterile water were more effective than "placebo" injections for relieving first-stage labor pain, 61

Inpatients

- Review: Sensitive thyrotropin testing in unselected inpatients has low diagnostic accuracy, 33

Internet

- Identifying systematic reviews: key resources (Editorial), A-11(No. 3)
- Netting the Evidence (Resource Corner), A-16(No. 1)

Intervertebral disk

- Review: Evidence supports surgery for lumbar disc prolapse but is insufficient for degenerative lumbar spondylosis, 100

Ipratropium

- Review: β -agonists plus ipratropium bromide improve airflow obstruction in adults with acute asthma, 12

K

Kidney disease

- Ramipril reduced the rate of progression to end-stage renal disease and overt proteinuria in nondiabetic nephropathy, 18
- Plasma volume expansion with albumin reduced renal impairment and death in cirrhosis and spontaneous bacterial peritonitis, 20

Knee joint

- Hyaluronic acid improved walking ability and short-term knee function in osteoarthritis of the knee, 98

L

Labor stage, first

- Cutaneous lumbosacral injections of sterile water were more effective than "placebo" injections for relieving first-stage labor pain, 61

Laser surgery

- Transmyocardial revascularization increased exercise duration and reduced angina symptoms, 85
- Transmyocardial revascularization was effective in refractory angina and left ventricular free-wall ischemia, 84

Lipoproteins, HDL cholesterol

- Gemfibrozil reduced the risk for coronary events in men with coronary artery disease and low levels of high-density lipoprotein cholesterol, 44

Lithium

- Review: Lithium augmentation increases treatment response in refractory depression, 105
- Review: Studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence, 26

Liver cirrhosis, biliary

- Review: Long-term ursodeoxycholic acid does not prevent mortality or morbidity in primary biliary cirrhosis, 68

Low back pain

- An 8-session exercise program was effective for subacute or chronic low-back pain, 99
- Review: The effectiveness of acupuncture for nonspecific low-back pain is unclear, 23

Lung diseases, obstructive

- Glucocorticoids reduced short-term treatment failure in exacerbations of chronic obstructive pulmonary disease, 14

Lung neoplasms

- Review: Transthoracic needle aspiration biopsy is accurate for diagnosing malignant and specific benign pulmonary lesions, 111

M

Mammography

- Review: Routine follow-up tests do not improve survival and quality of life in women after treatment for early-stage breast cancer, 106

Masks

- Therapeutic NCPAP was more effective than subtherapeutic NCPAP in obstructive sleep apnea, 16

Mastectomy, segmental

- Tamoxifen added to lumpectomy and radiation therapy reduced breast cancer events in ductal carcinoma in situ, 54

Maxillary sinusitis

- Review: Penicillin V or amoxicillin is better than placebo and equal to nonpenicillins for acute maxillary sinusitis, 47

Medical history taking

- Review: The physical examination can exclude the diagnosis of meningitis in low-risk adults, 32

Medline

- PubMed and SUMSearch (Resource Corner), A-16 (No. 3)

Meningitis

- Review: The physical examination can exclude the diagnosis of meningitis in low-risk adults, 32

Menstruation disorders

- Transcervical resection of the endometrium was effective and acceptable at up to 2 years for heavy menstrual bleeding, 28

Methylprednisolone

- Methylprednisolone injection improved symptoms for 1 year in patients with the carpal tunnel syndrome, 56

- Metoprolol
Controlled- and extended-release metoprolol reduced mortality in congestive heart failure, 3
- Midazolam
Buccal midazolam may be a useful alternative to rectal diazepam for treating acute seizures in refractory epilepsy, 22
- Mortality
Controlled- and extended-release metoprolol reduced mortality in congestive heart failure, 3
Increased time from assessment to angioplasty was associated with an increased 30-day mortality rate, 71
- Myocardial infarction
Dalteparin reduced death and MI at 1 but not 3 months in unstable coronary artery disease managed noninvasively, 82
Immediate revascularization for cardiogenic shock after MI reduced 6-month mortality, 42
Increased time from assessment to angioplasty was associated with an increased 30-day mortality rate, 71
Low-dose aspirin reduced combined 3-month risk for stroke, MI, and death after carotid endarterectomy, 10
n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction, 6
Primary angioplasty had lower long-term mortality than did streptokinase in acute myocardial infarction, 81
Review: Long-term β -blocker use reduces mortality after myocardial infarction, 4
Younger women with acute MI had more in-hospital deaths than men of the same age, 36
- Myocardial revascularization
Immediate revascularization for cardiogenic shock after MI reduced 6-month mortality, 42
An invasive strategy reduced the combined rate of death and myocardial infarction in unstable coronary artery disease, 83
Transmyocardial revascularization increased exercise duration and reduced angina symptoms, 85
Transmyocardial revascularization was effective in refractory angina and left ventricular free-wall ischemia, 84

N

- Nails, ingrown
Review: Avulsion with phenol prevents recurrence of ingrown toenails, 30
- Naltrexone
Review: Studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence, 26
- Nebulizers and vaporizers
Spacers made from sealed cold-drink bottles were as effective as conventional spacers for children with acute asthma, 102
- Neostigmine
Neostigmine improved acute colonic pseudo-obstruction quickly and effectively, 19
- Neutropenia
Empirical oral and IV antibiotics had similar effects for febrile neutropenia during chemotherapy in low-risk patients with cancer, 52,53

O

- Omeprazole
Omeprazole was better than cisapride or placebo for controlling heartburn at 4 and 8 weeks, 64

- Oseltamivir
Oseltamivir once or twice daily safely prevented influenza, 92
- Osteoarthritis
Celecoxib was similar to naproxen for rheumatoid arthritis with fewer endoscopic ulcers, 97
Hyaluronic acid improved walking ability and short-term knee function in osteoarthritis of the knee, 98
Rofecoxib caused fewer endoscopic gastroduodenal ulcers than ibuprofen in osteoarthritis, 96
Review: Exercise therapy may reduce pain and disability in osteoarthritis of the hip or knee, 62
- Osteoporosis
Raloxifene reduced vertebral fractures in postmenopausal women, 58
Review: External hip protectors reduce the risk for hip fractures in elderly persons, 63
Review: Vitamin D plus calcium is more effective than no therapy or calcium alone in corticosteroid-induced osteoporosis, 57
- Outcome assessment (health care)
Review: Interactive, but not didactic, continuing medical education is effective in changing physician performance, 75
- Outpatient clinics, hospital
Women with breast cancer were more satisfied with general practitioner care than with outpatient clinic care, 106

P

- Parents
The *Baby Check* booklet did not affect frequency of health service use in infants ≤ 6 months of age, 31
- Paroxetine
Paroxetine was effective for reducing symptoms in social phobia, 103
- Patient discharge
Organ system function, general health, social support, and age predicted discharge from a postacute geriatric unit to the community, 77
- Patient education
Attendance at an asthma clinic did not reduce asthma morbidity more than did standard medical treatment, 51
Review: Multiple interventions modestly reduce cardiovascular risk factors in primary prevention, but effects on mortality are uncertain, 43
- Patient readmission
A multidisciplinary, home-based intervention reduced deaths and readmissions in patients with chronic congestive heart failure, 88
- Patient satisfaction
Review: Health care decision aids improve knowledge, decrease decisional conflict, and increase active participation, 66
- Penicillins
Review: Penicillin V or amoxicillin is better than placebo and equal to nonpenicillins for acute maxillary sinusitis, 47
- Pergolide
Pergolide reduced involuntary periodic leg movements and improved sleep in the restless legs syndrome, 24
- Periodicals
Bigger and better (Editorial), A-11(No. 1)
- Pertussis vaccine
Review: Acellular vaccines prevent pertussis as effectively as whole-cell vaccines with fewer adverse events, 27

- Phenol
Review: Avulsion with phenol prevents recurrence of ingrown toenails, 30
- Phobic disorders
Paroxetine was effective for reducing symptoms in social phobia, 103
- Physical examination
Review: The physical examination can exclude the diagnosis of meningitis in low-risk adults, 32
Review: Routine follow-up tests do not improve survival and quality of life in women after treatment for early-stage breast cancer, 106
- Physicians, family
Women with breast cancer were more satisfied with general practitioner care than with outpatient clinic care, 106
- Platelet aggregation inhibitors
Review: Antithrombotic agents prevent stroke in nonvalvular atrial fibrillation, 86
- Pneumonia
A semirecumbent body position produced a lower rate of nosocomial pneumonia than did a supine position in mechanically ventilated adults, 89
- Pollen
Grass-pollen immunotherapy for 3 to 4 years was still effective 3 years after being discontinued, 65
- Positive-pressure respiration
Therapeutic NCPAP was more effective than subtherapeutic NCPAP in obstructive sleep apnea, 16
- Predictive value of tests
The ^{13}C -urea blood test is accurate for detecting *Helicobacter pylori* infection, 35
Whole-blood antibody tests were not highly sensitive for detecting *Helicobacter pylori* infection, 34
- Prednisone
Inhaled budesonide plus oral prednisone reduced relapse after discharge from the emergency department for acute asthma, 13
- Pregnancy
The high-risk criteria of a clinical prediction model were specific but not sensitive for predicting ectopic pregnancy, 117
Review: Antihypertensive drugs improve maternal outcomes in mild chronic and pregnancy-induced hypertension (Letter), A-19(No. 2)
- Primary health care
Omeprazole was better than cisapride or placebo for controlling heartburn at 4 and 8 weeks, 64
- Prone position
A semirecumbent body position produced a lower rate of nosocomial pneumonia than did a supine position in mechanically ventilated adults, 89
- Propranolol
Endoscopic ligation was better than propranolol for prevention of bleeding from esophageal varices, 21
- Prostatic hyperplasia
Review: β -sitosterols improve urinary symptoms in the short term in men with benign prostatic hyperplasia, 94
- Protective devices
Review: External hip protectors reduce the risk for hip fractures in elderly persons, 63
Use of an impervious wound-edge protector decreased postoperative wound infection, 60
- Proteinuria
Ramipril reduced the rate of progression to end-stage renal disease and overt proteinuria in nondiabetic nephropathy, 18
- Publishing
Bigger and better (Editorial), A-11(No. 1)
- Pyroles
Atorvastatin was at least as effective as PTCA for reducing ischemic events in patients with stable coronary artery disease, 7
- ## Q
- Quality of life
Empirical eradication therapy was better than endoscopy-based management for *Helicobacter pylori* infection and dyspepsia, 46
- ## R
- Ramipril
Ramipril reduced mortality and cardiovascular morbidity in high-risk adults, 41
Ramipril reduced the rate of progression to end-stage renal disease and overt proteinuria in nondiabetic nephropathy, 18
- Random allocation
Assessing allocation concealment and blinding in randomized controlled trials: why bother? (Editorial), A-11(No. 2)
Incorporating allocation concealment and blinding in randomized controlled trials (Editorial), A-13(No. 2)
- Reagent kits, diagnostic
Whole-blood antibody tests were not highly sensitive for detecting *Helicobacter pylori* infection, 34
- Research methods
Incorporating allocation concealment and blinding in randomized controlled trials (Editorial), A-13(No. 2)
- Recurrence
Antidepressants plus cognitive therapy reduced relapse in residual depression, 104
n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction, 6
Neostigmine improved acute colonic pseudo-obstruction quickly and effectively, 19
- Restless legs syndrome
Pergolide reduced involuntary periodic leg movements and improved sleep in the restless legs syndrome, 24
- Risk factors
Diastolic blood pressure and weight gain predicted future risk for hypertension, 37
A simple index predicted cardiac complications in elective major noncardiac surgery, 76
- ## S
- Sensitivity and specificity
Review: Sensitive thyrotropin testing in unselected inpatients has low diagnostic accuracy, 33
The ^{13}C -urea blood test is accurate for detecting *Helicobacter pylori* infection, 35
Whole-blood antibody tests were not highly sensitive for detecting *Helicobacter pylori* infection, 34
- Serotonin agents
Review: Studies with sufficient follow-up do not show a clear benefit for pharmacotherapy in alcohol dependence, 26

Sex factors

Younger women with acute MI had more in-hospital deaths than men of the same age, 36

Shock, cardiogenic

Immediate revascularization for cardiogenic shock after MI reduced 6-month mortality, 42

Single-blind method

Assessing allocation concealment and blinding in randomized controlled trials: why bother? (Editorial), A-11 (No. 2)

Sitosterols

Review: β -sitosterols improve urinary symptoms in the short term in men with benign prostatic hyperplasia, 94

Skilled nursing facilities

Organ system function, general health, social support, and age predicted discharge from a postacute geriatric unit to the community, 77

Skin neoplasms

Daily use of β -carotene supplements did not prevent skin cancer, but daily sunscreen use reduced the incidence of squamous-cell carcinomas, 101

Sleep apnea syndrome

Review: Screening tests are not as accurate as overnight polysomnography for the diagnosis of adult sleep apnea, 69
Therapeutic NCPAP was more effective than subtherapeutic NCPAP in obstructive sleep apnea, 16

Sleep

Pergolide reduced involuntary periodic leg movements and improved sleep in the restless legs syndrome, 24

Software

Advances in evidence-based information resources for clinical practice (Editorial), A-11(No. 1)

Spinal fractures

Raloxifene reduced vertebral fractures in postmenopausal women, 58

Spironolactone

Spironolactone reduced mortality in severe congestive heart failure, 2

Stents

Abciximab and stenting reduced death, myocardial infarction, and repeated revascularization in coronary revascularization, 5

Stomach neoplasms

Review: Pooled heterogeneous studies show an association between *Helicobacter pylori* infection and gastric cancer, 74

Streptokinase

Primary angioplasty had lower long-term mortality than did streptokinase in acute myocardial infarction, 81

Sunscreening agents

Daily use of β -carotene supplements did not prevent skin cancer, but daily sunscreen use reduced the incidence of squamous-cell carcinomas, 101

Surgery

Endoscopic ligation was better than propranolol for prevention of bleeding from esophageal varices, 21

Laparoscopic hernia repair reduced recovery time and days of postoperative analgesia use better than Shouldice repair, 29

A simple index predicted cardiac complications in elective major noncardiac surgery, 76

Surgical volume was not related to 30-day mortality in 8 common operations, 72

Use of an impervious wound-edge protector decreased postoperative wound infection, 60

T

Tamoxifen

Tamoxifen added to lumpectomy and radiation therapy reduced breast cancer events in ductal carcinoma in situ, 54

Tedelparin

Dalteparin reduced death and MI at 1 but not 3 months in unstable coronary artery disease managed noninvasively, 82

Thyroid diseases

Review: Sensitive thyrotropin testing in unselected inpatients has low diagnostic accuracy, 33

Thyroid function tests

Review: Sensitive thyrotropin testing in unselected inpatients has low diagnostic accuracy, 33

Tissue plasminogen activators

Tissue plasminogen activator improved function at 6 and 12 months after ischemic stroke, 11

Tomography, x-ray computed

Virtual colonoscopy was sensitive and specific for detecting colorectal polyps and cancer, 110

Trimethoprim-sulfamethoxazole combination

Trimethoprim-sulfamethoxazole decreased morbidity and mortality in HIV-1-infected patients with tuberculosis, 45

Tuberculosis

Trimethoprim-sulfamethoxazole decreased morbidity and mortality in HIV-1-infected patients with tuberculosis, 45

U

Ursodeoxycholic acid

Review: Long-term ursodeoxycholic acid does not prevent mortality or morbidity in primary biliary cirrhosis, 68

V

Vegetables

Fruit and vegetable intake decreased risk for ischemic stroke, 115

Venous thrombosis

A negative D-dimer result and low-risk clinical status effectively ruled out DVT in symptomatic patients, 108

A negative D-dimer test result alone or combined with low-risk clinical status effectively ruled out symptomatic DVT, 109

Veterans

Integrated outpatient treatment increased abstinence in men with alcohol-related illness who were ongoing drinkers, 67

Vitamin B complex

Review: High levels of homocysteine are associated with an increased risk for cardiovascular disease, 73

Vitamin D

Review: Vitamin D plus calcium is more effective than no therapy or calcium alone in corticosteroid-induced osteoporosis, 57

Vitamin E

n-3 polyunsaturated fatty acids reduced mortality and morbidity after recent myocardial infarction, 6

W

Whooping cough

Review: Acellular vaccines prevent pertussis as effectively as whole-cell vaccines with fewer adverse events, 27

Z

Zanamivir

Zanamivir speeds symptom relief in influenza, 93

